CTC2, NWI-MOL176, exercises week 2

Gerrit C. Groenenboom, 19-April-2023

Question 1: Questions chapter 5

In section 5.1 of the lecture notes, the angular momentum states $|ab\rangle$ are defined by

$$\hat{l}^2 |ab\rangle = a\hbar^2 |ab\rangle \tag{1}$$

$$\hat{l}_z |ab\rangle = b\hbar |ab\rangle. \tag{2}$$

Ladder operators are defined by

$$\hat{l}_{\pm} = \hat{l}_x \pm i \hat{l}_y. \tag{3}$$

1a. Show that $b^2 \leq a$, using Eqs. (5.1)-(5.4) of the lecture notes and the properties of scalar products (section 1.5.3 of the lecture notes)

Answer: The expectation value of the square of a Hermitian operator must be positive, e.g.,

$$\langle ab|\hat{l}_x^2|ab\rangle = \langle \hat{l}_x ab||\hat{l}_x ab\rangle = |\hat{l}_x|ab\rangle|^2 \ge 0.$$
(4)

Since

$$\langle ab|\hat{l}^2|ab\rangle - \langle ab|\hat{l}^2_z|ab\rangle i = \langle ab|\hat{l}^2_x + \hat{l}^2_y + \hat{l}^2_z|ab\rangle - \langle ab|\hat{l}^2_z|ab\rangle = \langle ab|\hat{l}^2_x|ab\rangle + \langle ab|\hat{l}^2_y|ab\rangle \ge 0$$

$$a\hbar^2 - (b\hbar)^2 \ge 0$$
(5)

$$(6)$$

$$a \ge b^2. \tag{7}$$

1b. The angular momentum ladder operators are each other's Hermitian conjugates, $\hat{l}^{\dagger}_{\pm} = \hat{l}_{\pm}$. Derive this result using the definition of Hermitian conjugate and the defining properties of scalar products.

Answer:

$$\hat{l}_{\pm} = \hat{l}_x \pm i \hat{l}_y. \tag{8}$$

Since \hat{l}_x is Hermitian we have $\hat{l}_x^{\dagger} = \hat{l}_x$. Thus, we have to show

$$(i\hat{l}_y)^{\dagger} = -\hat{l}_y. \tag{9}$$

Since \hat{l}_y is Hermitian, for any state $s|\phi\rangle$, χ we have

$$\langle \phi | \hat{l}_y \chi \rangle = \langle \hat{l}_y \phi | \chi \rangle \tag{10}$$

Thus, for these states we also have,

$$\langle \phi | i \hat{l}_y \chi \rangle = i \langle \hat{l}_y \phi | \chi \rangle = \langle -i \hat{l}_y \phi | \chi \rangle, \tag{11}$$

and hence $(i\hat{l}_y)^{\dagger} = -i\hat{l}_y$, q.e.d.

1c. Show that

$$\hat{l}_{\pm}\hat{l}_{\mp} = \hat{l}^2 - \hat{l}_z^2 \pm \hbar \hat{l}_z.$$

Answer:

$$\hat{l}_{\pm}\hat{l}_{\mp} = (\hat{l}_x \pm i\hat{l}_y)(\hat{l}_x \mp i\hat{l}_y)$$
 (12)

$$=\hat{l}_{x}^{2}+\hat{l}_{y}^{2}\mp i[\hat{l}_{x},\hat{l}_{y}]$$
(13)

- (14)
- $= \hat{l}_x^2 + \hat{l}_y^2 \mp i^2 \hbar \hat{l}_z$ $= \hat{l}^2 \hat{l}_z^2 \pm \hbar \hat{l}_z.$ (15)

1d. Compute the matrix representation of \hat{l}^2 , \hat{l}_z , \hat{l}_+ , \hat{l}_- , \hat{l}_x , and \hat{l}_y in the angular momentum basis $\{|lm\rangle, m = -l, -l+1, \dots, l\}$, with l = 1.

Answer:

$$L^{2} = \hbar^{2} \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$$
$$L_{z} = \hbar \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
$$L_{+} = \hbar \begin{pmatrix} 0 & 0 & 0 \\ \sqrt{2} & 0 & 0 \\ 0 & \sqrt{2} & 0 \end{pmatrix}$$
$$L_{-} = \hbar \begin{pmatrix} 0 & \sqrt{2} & 0 \\ 0 & 0 & \sqrt{2} \\ 0 & 0 & 0 \end{pmatrix}$$
$$L_{x} = \frac{\hbar}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

- 1e. Check the relation $\hat{l}^2 = \hat{l}_z^2 + \hbar \hat{l}_z + \hat{l}_- \hat{l}_+$ for the l = 1 matrix representations from the previous question.
- 1f. Compute

for the 2×2 matrix

$$\boldsymbol{A} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

 $\boldsymbol{B}=\cos(\boldsymbol{A}),$

Hint: first solve the eigenvalue problem, Eq. (5.35) in the lecture notes and then use Eq. (5.47) with $f(\lambda_i) = \cos(\lambda_i)$.

According to section 5.6 of the lecture notes, the translation operator [Eq. (5.54)] acting on $\sin(x)$ gives [Eq. (5.49)]

$$e^{-\alpha \frac{\partial}{\partial x}} \sin(x) = \sin(x - \alpha).$$

1g. Show that this equation is correct in first order, i.e., for small α .

1h. Compute

$$\left(e^{-3\frac{\partial}{\partial x}}\right)^2 e^{-x^2}.$$

1i. What is wrong in this derivation:

$$e^{\frac{\partial}{\partial x}}e^x = e^{\frac{\partial}{\partial x}x} = e^1 = e?$$

1j. Repeat question 1f, but now for the complex matrix

$$\boldsymbol{A} = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}.$$

The angular momentum operator \hat{l}_z in spherical polar coordinates, is given by [see lecture notes Eq. (5.66)]

$$\hat{l}_z = \frac{\hbar}{i} \frac{\partial}{\partial \phi}.$$
(16)

1k. Derive this result starting from the expression for \hat{l}_z in Cartesian coordinates.

Answer: From the definition:

$$\hat{l}_z = x\hat{p}_y - y\hat{p}_x = \frac{\hbar}{i} \left(x\frac{\partial}{\partial y} - y\frac{\partial}{\partial x} \right).$$
(17)

With

$$x = r\cos\phi\sin\theta \tag{18}$$

$$y = r\sin\phi\sin\theta \tag{19}$$

$$z = r\cos\theta \tag{20}$$

and starting from Eq. (16) we find

$$\hat{l}_z = \frac{\hbar}{i} \frac{\partial}{\partial \phi} \tag{21}$$

$$=\frac{\hbar}{i}\left(\frac{\partial y}{\partial \phi}\frac{\partial}{\partial y}+\frac{\partial x}{\partial \phi}\frac{\partial}{\partial x}+\frac{\partial z}{\partial \phi}\frac{\partial}{\partial z}\right)$$
(22)

$$=\frac{\hbar}{i}\left(r\cos\phi\sin\theta\frac{\partial}{\partial y} - r\sin\phi\sin\theta\frac{\partial}{\partial x} + 0\right)$$
(23)

$$=\frac{\hbar}{i}\left(x\frac{\partial}{\partial y}-y\frac{\partial}{\partial x}\right),\tag{24}$$

q.e.d.