

# Ocean-like water in the Jupiter-family comet 103P/Hartley 2

Paul Hartogh<sup>1</sup>, Dariusz C. Lis<sup>2</sup>, Dominique Bockelée-Morvan<sup>3</sup>, Miguel de Val-Borro<sup>1</sup>, Nicolas Biver<sup>3</sup>, Michael Küppers<sup>4</sup>, Martin Emprechtinger<sup>2</sup>, Edwin A. Bergin<sup>5</sup>, Jacques Crovisier<sup>3</sup>, Miriam Rengel<sup>1</sup>, Raphael Moreno<sup>3</sup>, Slawomira Szutowicz<sup>6</sup> & Geoffrey A. Blake<sup>2</sup>


For decades, the source of Earth's volatiles, especially water with a deuterium-to-hydrogen ratio (D/H) of  $(1.558 \pm 0.001) \times 10^{-4}$ , has been a subject of debate. The similarity of Earth's bulk composition to that of meteorites known as enstatite chondrites<sup>1</sup> suggests a dry proto-Earth<sup>2</sup> with subsequent delivery of volatiles<sup>3</sup> by local accretion<sup>4</sup> or impacts of asteroids or comets<sup>5,6</sup>. Previous measurements in six comets from the Oort cloud yielded a mean D/H ratio of  $(2.96 \pm 0.25) \times 10^{-4}$ . The D/H value in carbonaceous chondrites,  $(1.4 \pm 0.1) \times 10^{-4}$ , together with dynamical simulations, led to models in which asteroids were the main source of Earth's water<sup>7</sup>, with  $\leq 10$  per cent being delivered by comets. Here we report that the D/H ratio in the Jupiter-family comet 103P/Hartley 2, which originated in the Kuiper belt, is  $(1.61 \pm 0.24) \times 10^{-4}$ . This result substantially expands the reservoir of Earth ocean-like water to include some comets, and is consistent with the emerging picture of a complex dynamical evolution of the early Solar System<sup>8,9</sup>.

On 17 November 2010, using the Herschel Space Observatory, we determined the D/H ratio in a comet from a reservoir other than the Oort cloud—103P/Hartley 2. Such Jupiter-family comets are believed to originate from the Kuiper belt, which exists beyond the orbits of the giant planets at radii between 30 and 50 astronomical units<sup>10</sup> (1 AU is the average Earth–Sun distance). In contrast, Oort-cloud comets are theorized to have originated from radii near the gas giants and to have been subsequently ejected to the Oort cloud ( $>5,000$  AU)<sup>11</sup>. The Herschel measurement therefore traces the water D/H ratio in a new population of water-ice-rich bodies in the Solar System that are a potential source of water on the Earth.

To obtain an accurate determination of the D/H ratio in water, we carried out simultaneous observations of optically thin isotopic variants of water, specifically HDO and H<sub>2</sub><sup>18</sup>O (Fig. 1), as part of our Solar System observing programme<sup>12</sup>. This was critical for comet 103P/Hartley 2, whose activity and water outgassing rates exhibited significant short-term variations<sup>13</sup>. We used state-of-the-art excitation models to determine the HDO and H<sub>2</sub><sup>18</sup>O beam integrated column densities and production rates from the measured line intensities. Observation and modelling details are given in Supplementary Information. A critical point is that all observations sampled the same region of the coma, about 6,500 km in diameter.

The retrieved gas column densities and production rates are sensitive to collisional cross-sections, along with the density and temperature profiles of H<sub>2</sub>O and electrons, and we thus considered a range of model parameters (Table 1). Although the production rates determined for the various model parameters differ slightly, the value of the D/H ratio is estimated to be  $(1.61 \pm 0.24) \times 10^{-4}$ . In our analysis, we assumed an H<sub>2</sub><sup>16</sup>O/H<sub>2</sub><sup>18</sup>O ratio of  $500 \pm 50$ , a range that encompasses the Earth value and is consistent with previous measurements in cometary water<sup>14</sup> (see also Supplementary Information). The quoted  $1\sigma$  uncertainty in the D/H ratio includes a 5% uncertainty related to modelling.

Our measured D/H value is substantially larger than that which characterized the young Sun (4.5 Gyr ago; the protosolar ratio), believed to be about  $2.1 \times 10^{-5}$ , which in turn is slightly higher than the value found in the local interstellar medium today ( $1.6 \times 10^{-5}$ ) and



**Figure 1 | Submillimetre water emission lines from comet 103P/Hartley 2.** The time of the observations was 20 days after perihelion, when the comet was 1.095 AU from the Sun and 0.212 AU from Herschel. Because the H<sub>2</sub>O ground state rotational lines in comets are optically thick<sup>29,30</sup>, observations of the rare oxygen isotopic counterpart, H<sub>2</sub><sup>18</sup>O, provide a more reliable reference for the D/H determination. The spectra of the  $1_{10}-1_{01}$  lines of HDO (a) and H<sub>2</sub><sup>18</sup>O (b) at 509.292 and 547.676 GHz, respectively, were obtained with the Heterodyne Instrument for the Far Infrared (HIFI) High Resolution Spectrometer (HRS) between 17.28 and 17.64 November 2010 UT. The line intensities, expressed in the main-beam brightness temperature scale, are  $0.011 \pm 0.001$  and  $0.117 \pm 0.002$  K km s<sup>-1</sup>, for HDO and H<sub>2</sub><sup>18</sup>O respectively, averaging the two instrument polarizations. The velocity scale is given relative to the velocity of the comet's nucleus. The spectral resolution is 141 and 132 m s<sup>-1</sup> for the HDO and H<sub>2</sub><sup>18</sup>O spectra, respectively. For details of the observational sequence and basic parameters of the data analysis, see Supplementary information.

<sup>1</sup>Max-Planck-Institut für Sonnensystemforschung, Max-Planck-Str. 2, 37191 Katlenburg-Lindau, Germany. <sup>2</sup>California Institute of Technology, Pasadena, California 91125, USA. <sup>3</sup>LESIA-Observatoire de Paris, CNRS, UPMC, Université Paris-Diderot, 5 place Jules Janssen, 92195 Meudon, France. <sup>4</sup>Rosetta Science Operations Centre, European Space Astronomy Centre, 28691 Villanueva de la Cañada, Madrid, Spain. <sup>5</sup>Astronomy Department, University of Michigan, Ann Arbor, Michigan 48109, USA. <sup>6</sup>Space Research Centre, Polish Academy of Sciences, 00-716 Warsaw, Poland.

**Table 1 | Calculating the D/H ratio in water in comet 103P/Hartley 2**

Model	$T_{\text{gas}}$ (K)	$x_{\text{ne}}$	$\langle N(\text{HDO}) \rangle$ ( $10^{10} \text{ cm}^{-2}$ )	$Q(\text{HDO})$ ( $10^{24} \text{ s}^{-1}$ )	$\langle N(\text{H}_2^{18}\text{O}) \rangle$ ( $10^{11} \text{ cm}^{-2}$ )	$Q(\text{H}_2^{18}\text{O})$ ( $10^{25} \text{ s}^{-1}$ )	D/H
(1)	50	0.	4.9	3.1	3.5	2.1	$1.49 \times 10^{-4}$
	50	0.2	3.6	2.3	2.5	1.5	$1.55 \times 10^{-4}$
	70	0.2	3.7	2.4	2.5	1.5	$1.60 \times 10^{-4}$
	Law	0.1	5.7	3.6	3.7	2.2	$1.63 \times 10^{-4}$
(2)	50	0.2	4.3	2.7	2.9	1.8	$1.54 \times 10^{-4}$
	Law	0.1	4.8	3.1	3.2	1.9	$1.58 \times 10^{-4}$


The parameter  $x_{\text{ne}}$ , scaling the electron density profile in the models, is constrained by mapping observations.  $\langle N \rangle$  and  $Q$  are respectively beam integrated column density and production rate, determined using different parameters in the excitation models<sup>14,29</sup>. Production rates were computed assuming isotropic outflow of water from the nucleus, with a velocity of  $0.6 \text{ km s}^{-1}$ , consistent with the width of the  $\text{H}_2^{18}\text{O}$  line. We accounted for the  $10''$  offset between the centre of the beam and the position of the peak of the  $\text{H}_2\text{O}$  distribution. Values of 50 and 70 K for the gas kinetic temperature,  $T_{\text{gas}}$ , are consistent with multi-transition measurements of gaseous species in the millimetre and near-infrared range, respectively. The gas kinetic temperature is expected to decrease with increasing distance owing to quasi-adiabatic expansion of the escaping gases: the temperature law assumes that  $T_{\text{gas}} = 80 \text{ K}$  for  $r < 270 \text{ km}$ ,  $T_{\text{gas}} = 12 \text{ K}$  for  $r > 630 \text{ km}$ , with a linear decrease between 270 and 630 km, where  $r$  is the distance from the nucleus. Collision cross-sections involving water molecules and electrons are modelled differently in models (1) and (2). Both models use an electron density profile based on *in situ* measurements in comet 1P/Halley scaled to the activity of 103P/Hartley 2 (ref. 29). The D/H ratio is equal to  $0.5 \times Q(\text{HDO})/Q(\text{H}_2\text{O})$ , with  $Q(\text{H}_2\text{O}) = 500 \times Q(\text{H}_2^{18}\text{O})$ . See Supplementary information for details of the models and model parameters.

comparable to the primordial D/H ratio in the Universe after the Big Bang (Fig. 2)<sup>15</sup>. Protosolar water, on the other hand, is believed to be highly enriched (D/H  $\approx 1 \times 10^{-3}$ )<sup>16</sup> due to the low-temperature ( $\sim 10$ – $30 \text{ K}$ ) non-equilibrium chemistry that characterizes the dense interstellar medium<sup>17</sup>, either via gas-phase isotopic exchange reactions involving ions and radicals, or grain-surface processes. Consequently, the resulting D/H ratio in water ice is very sensitive to the physical conditions, in particular the kinetic temperature of the medium. After the protosolar cloud collapsed to form the solar nebula, isotopic exchange reactions between molecular hydrogen and HDO molecules would have led to a gradual reduction of D/H in water<sup>18</sup>, as compared to the initial interstellar medium value. Because the efficiency of these reactions and the turbulent mixing within the solar nebula is correlated with the gas density and temperature, the deuterium enhancement in water has been predicted to increase with the heliocentric distance<sup>19–21</sup>. Ices, captured by planetesimals and cometesimals, would have then preserved the deuterium enrichment in water from this early epoch. As

a result, small Solar System bodies are expected to exhibit different D/H ratios in their water ice depending on the distance from the Sun at which they were formed.

In the context of this simple nebular model, the D/H ratio of ( $1.61 \pm 0.24$ )  $\times 10^{-4}$  in comet 103P/Hartley 2—a factor of two lower than that measured in Oort-cloud comets (Fig. 2) and, within uncertainties, consistent with that of the Earth's oceans (for which the Vienna Standard Mean Ocean Water (VSMOW) value is ( $1.558 \pm 0.001$ )  $\times 10^{-4}$ )—is therefore surprising, and compatible with two different schemes: (1) either this comet did not form in a region that was further from the Sun than the assembly zone of the Oort-cloud comets, or (2) the dependence of the water D/H ratio with distance from the Sun is not as expected on the basis of current models. Concerning the first possibility, dynamical models indeed suggest that a fraction of the Jupiter-family comets originate in the Oort cloud<sup>22</sup>. Still, even if comet 103P/Hartley 2 stems from the Oort cloud, this would not explain why its D/H ratio is different from that seen in other Oort-cloud comets. Models also suggest that a fraction of the Jupiter-family comets may have originated from the Trojan asteroid swarms sharing the orbit of Jupiter<sup>23</sup>. The Trojans are generally thought to have resided at their current location since the formation of the Solar System. Therefore, Jupiter family comets originating in the Trojan region could, in theory, display deuterium enrichment values lower than those for bodies originating in the Kuiper belt, if they indeed formed in the vicinity of Jupiter. However, the most probable scenario is that 103P/Hartley 2 originated in the Kuiper belt.

It is difficult to explain the low D/H ratio in 103P/Hartley 2 (compared to that of previous measurements in comets) with the formation regions of comets, thus models of the gradient of D/H in the Solar System—predictions not yet directly confirmed by observations, owing to scarcity of accurate isotopic measurements—may need to be revisited. In fact, one recent model has suggested that the D/H ratio of water vapour can be locally enhanced<sup>24</sup>. However, the vapour must then be implanted into cometary ices. Moreover, until the measurement of 103P/Hartley 2 there was no observational confirmation of variations in the D/H ratio. One possible solution is that there was large-scale movement of material between the inner and outer Solar



**Figure 2 | D/H ratios in the Solar System.** Orange squares, values measured for water in the Oort-cloud comets 1P/Halley, C/1996 B2 (Hyakutake), C/1995 O1 (Hale-Bopp), C/2002 T7 (LINEAR) and 8P/Tuttle. Arrow (for 153P/Ikeya-Zhang), upper limit. Purple square, present measurement in the water of 103P/Hartley 2. Black symbols, D/H ratio in  $\text{H}_2$  in the atmosphere of the giant planets—Jupiter (J), Saturn (S), Uranus (U) and Neptune (N). Light blue and green symbols, D/H values for water in the plume of Saturn's moon Enceladus and in CI carbonaceous chondrites, respectively. Error bars,  $1\sigma$ . The D/H

determinations in comets originating from the Oort cloud are twice the value for the Earth's ocean (blue line) and about a factor of ten larger than the protosolar value in  $\text{H}_2$  (broad yellow line), the latter being comparable to the value in atomic hydrogen found in the local interstellar medium (ISM, red horizontal line). The D/H ratio in the Jupiter-family comet 103P/Hartley 2 is the same as the Earth's ocean value and the chondritic CI value. Uranus and Neptune have been enriched in deuterium by the mixing of their atmospheres with D-rich protoplanetary ices. For further details, see Supplementary Table 1.

System. According to a recent theory proposed for the early Solar System (the Grand Tack scenario), when the giant planets were still embedded in the nebular gas disk, there was a general radial mixing of the distribution of comets and asteroids born in different regions<sup>25</sup>. The similarity of the D/H ratio in comet 103P/Hartley 2, which probes the Kuiper belt, with that found in CI chondrites tracing the asteroid belt, would be in agreement with a general shake-up of the Solar System at early times.

In more 'orderly' models, the high D/H values derived from the earlier observations of Oort-cloud comets suggested that at most 10% of the Earth's water could have been supplied from the outermost Solar System, but even under these circumstances a number of scenarios have been developed suggesting that terrestrial water could have in fact been delivered by comets. Such models are based on assumptions about the heliocentric D/H gradient<sup>26</sup>, and the analysis of lunar samples<sup>27</sup> and telluric sedimentary rocks formed at the end of the Late Heavy Bombardment phase<sup>28</sup>. Our Herschel observations of a VSMOW-like D/H ratio in 103P/Hartley 2 enlarge the region of the solar nebula known to have a D/H ratio similar to that of Earth's oceans; this region now includes both the asteroid belt and the much larger Kuiper belt, thereby providing support for the theory of a common water source for the inner Solar System bodies (including the Earth) in which comets play an important part.

Further constraints on the delivery of volatiles to the early Earth and an improved understanding of the origin of the different dynamical classes of comets will require significantly larger sample sizes than those at present available. A handful of additional measurements can be expected from Herschel before its cryogen supply is exhausted, but the comparison of D/H ratios in the inner and outer Solar System must necessarily utilize very different objects and materials. For the inner Solar System, *in situ* space missions or sample return missions to the outer asteroid belt would provide critical new data. Astronomically, the minuscule strength of HDO spectroscopic signatures makes D/H measurements extremely challenging, and dedicated programmes using new facilities will be required to substantially increase the inventory of high-precision D/H ratios in comets and other icy Solar System bodies, including the Jovian satellites.

Received 16 May; accepted 30 August 2011.

Published online 5 October 2011.

- Javoy, M. *et al.* The chemical composition of the Earth: enstatite chondrite models. *Earth Planet. Sci. Lett.* **293**, 259–268 (2010).
- Wänke, H. Constitution of terrestrial planets. *Phil. Trans. R. Soc. Lond.* **303**, 287–302 (1981).
- Robert, F. A distinct source of lunar water? *Nature Geosci.* **4**, 74–75 (2011).
- Drake, M. J. & Righter, K. Determining the composition of the Earth. *Nature* **416**, 39–44 (2002).
- Oró, J. Comets and the formation of biochemical compounds on the primitive Earth. *Nature* **190**, 389–390 (1961).
- Owen, T., Bar-Nun, A. & Kleinfeld, I. Possible cometary origin of heavy noble gases in the atmospheres of Venus, Earth and Mars. *Nature* **358**, 43–46 (1992).
- Morbidelli, A. *et al.* Source regions and time scales for the delivery of water to Earth. *Meteorit. Planet. Sci.* **35**, 1309–1320 (2000).
- Tsiganis, K., Gomes, R., Morbidelli, A. & Levison, H. F. Origin of the orbital architecture of the giant planets of the Solar System. *Nature* **435**, 459–461 (2005).
- Gomes, R., Levison, H. F., Tsiganis, K. & Morbidelli, A. Origin of the cataclysmic Late Heavy Bombardment period of the terrestrial planets. *Nature* **435**, 466–469 (2005).
- Levison, H. F. & Duncan, M. J. From the Kuiper belt to Jupiter-family comets: the spatial distribution of ecliptic comets. *Icarus* **127**, 13–32 (1997).
- Dones, L., Weissman, P. R., Levison, H. F. & Duncan, M. J. in *Comets II* (eds Festou, M. C., Keller, H. U. & Weaver, H. A.) 153–174 (Univ. Arizona Press, 2005).
- Hartogh, P. *et al.* Water and related chemistry in the solar system: a guaranteed time key programme for Herschel. *Planet. Space Sci.* **57**, 1596–1606 (2009).

- Meech, K. *et al.* EPOXI: 103P/Hartley 2 observations from a worldwide campaign. *Astrophys. J.* **734**, L1 (2011).
- Biver, N. *et al.* Submillimetre observations of comets with Odin: 2001–2005. *Planet. Space Sci.* **55**, 1058–1068 (2007).
- Linsky, J. L. *et al.* What is the total deuterium abundance in the local galactic disk? *Astrophys. J.* **647**, 1106–1124 (2006).
- Butner, H. M. *et al.* Discovery of interstellar heavy water. *Astrophys. J.* **659**, L137–L140 (2007).
- Watson, W. D. Ion-molecule reactions, molecule formation, and hydrogen-isotope exchange in dense interstellar clouds. *Astrophys. J.* **188**, 35–42 (1974).
- Geiss, J. & Reeves, H. Cosmic and solar system abundances of deuterium and helium-3. *Astron. Astrophys.* **18**, 126–132 (1972).
- Drouart, A., Dubrulle, B., Gautier, D. & Robert, F. Structure and transport in the solar nebula from constraints on deuterium enrichment and giant planet formation. *Icarus* **140**, 129–155 (1999).
- Mousis, O. *et al.* Constraints on the formation of comets from D/H ratios measured in H<sub>2</sub>O and HCN. *Icarus* **148**, 513–525 (2000).
- Aikawa, Y. & Herbst, E. Two-dimensional distributions and column densities of gaseous models in protoplanetary disks II. Deuterated species and UV shielding by ambient clouds. *Astron. Astrophys.* **371**, 1107–1117 (2001).
- Emel'yanenko, V. V., Asher, D. J. & Bailey, M. E. Centaurs from the Oort cloud and the origin of Jupiter-family comets. *Mon. Not. R. Astron. Soc.* **361**, 1345–1351 (2005).
- Marzari, F., Farinella, P. & Vanzani, V. Are Trojan collisional families a source for short-period comets? *Astron. Astrophys.* **299**, 267–276 (1995).
- Thi, W.-F., Woitke, P. & Kamp, I. Warm non-equilibrium gas phase chemistry as a possible origin of high HDO/H<sub>2</sub>O ratios in hot and dense gases: application to inner protoplanetary discs. *Mon. Not. R. Astron. Soc.* **407**, 232–246 (2010).
- Walsh, K. J. & Morbidelli, A. The effect of an early planetesimal-driven migration of the giant planets on terrestrial planet formation. *Astron. Astrophys.* **526**, A126 (2011).
- Delsemme, A. H. The deuterium enrichment observed in recent comets is consistent with the cometary origin of seawater. *Planet. Space Sci.* **47**, 125–131 (1998).
- Greenwood, J. P. *et al.* Hydrogen isotope ratios in lunar rocks indicate delivery of cometary water to the Moon. *Nature Geosci.* **4**, 79–82 (2011).
- Jørgensen, U. G. *et al.* The Earth-Moon system during the late heavy bombardment period — geochemical support for impacts dominated by comets. *Icarus* **204**, 368–380 (2009).
- Zakharov, V., Bockelée-Morvan, D., Biver, N., Crovisier, J. & Lecacheux, A. Radiative transfer simulation of water rotational excitation in comets. Comparison of the Monte Carlo and escape probability methods. *Astron. Astrophys.* **473**, 303–310 (2007).
- Bensch, F. & Bergin, E. A. The pure rotational line emission of ortho-water in comets. I. Radiative transfer model. *Astrophys. J.* **615**, 531–544 (2004).

Supplementary Information is linked to the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

**Acknowledgements** Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation by NASA. The Heterodyne Instrument for the Far Infrared (HIFI) has been designed and built by a consortium of institutes and university departments from across Europe, Canada and the United States under the leadership of SRON, the Netherlands Institute for Space Research, and with major contributions from Germany, France and the USA. This development has been supported by national funding agencies: CEA, CNES, CNRS (France); ASI (Italy); and DLR (Germany). Additional funding support for some instrument activities has been provided by ESA. Support for this work was also provided by NASA through an award issued by JPL/Caltech. D.C.L. is supported by an NSF award to the Caltech Submillimeter Observatory. We thank R. Lorente, P. García-Lario, M. Kidger and G. Pilbratt for helping with the scheduling of these observations, and I. Avruch for the assistance with HIFI specific software issues.

**Author Contributions** This paper represents the combined work of the HsO (the Herschel guaranteed time key programme "Water and related chemistry in the solar system") team members listed as authors. P.H. is the coordinator of this programme. All authors contributed to this work, including observation planning, data analysis and writing of the manuscript. N.B., D.B.-M., M.R., R.M., M.d.V.-B. and M.E. carried out the data reduction and contributed to the modelling efforts. All authors were collectively involved in the discussion and interpretation of the results.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at [www.nature.com/nature](http://www.nature.com/nature). Correspondence and requests for materials should be addressed to P.H. ([hartogh@mps.mpg.de](mailto:hartogh@mps.mpg.de)).